

Ultraviolet divergences of flavor changing amplitudes in the littlest Higgs model with T-parity

Toru Goto^{a,1}, Yasuhiro Okada^{a,b,2} and Yasuhiro Yamamoto^{a,b,3}

^a*Theory Group, KEK, Tsukuba, Ibaraki 305-0801, Japan*

^b*Department of Particle and Nuclear Physics, The Graduate University for Advanced
Studies (Sokendai), Tsukuba, Ibaraki 305-0801, Japan*

September 27, 2008

Abstract

Flavor changing neutral current processes are studied in the littlest Higgs model with T-parity. It is found that the logarithmic divergence reported earlier in Z boson flavor changing processes is exactly canceled by contributions from additional interaction terms of heavy fermions and the Z boson. Phenomenological impact on the $K \rightarrow \pi \nu \bar{\nu}$ processes is discussed.

¹e-mail: tgoto@post.kek.jp

²e-mail: yasuihiro.okada@kek.jp

³e-mail: yamayasu@post.kek.jp

The little Higgs model [1, 2] was proposed as a solution to the little hierarchy problem [3] by canceling the quadratic divergence of the Higgs mass term at one-loop level due to new diagrams with additional gauge bosons and a heavy top-quark partner. It was soon realized that the scale of the new particles should be in the multi-TeV range in order to satisfy constraints from electroweak precision measurements, so that the little hierarchy problem is reintroduced [4]. This problem is avoided in the little Higgs model with T-parity [5], because dangerous diagrams with the tree level exchange of heavy neutral gauge bosons are forbidden by a new Z_2 discrete symmetry.

There are new sources of flavor transition in the littlest Higgs model with T-parity. In order to assign the Z_2 symmetry for quarks and leptons, new heavy fermions have to be introduced, and flavor mixing matrices associated with the heavy fermions are independent of the Cabibbo-Kobayashi-Maskawa (CKM) [6] or the Pontecorvo-Maki-Nakagawa-Sakata (PMNS) [7] matrices [8]. Loop diagrams with heavy fermions and gauge bosons can induce new contributions to quark flavor changing neutral current (FCNC) processes and lepton flavor violating (LFV) processes. In Refs. [9, 10, 11], these processes are studied in detail and large deviation from the Standard Model predictions are shown to be possible. In these works, it is argued that there are “leftover” singularities in FCNC/LFV amplitudes, and these logarithmic divergent terms can be dominant contributions.

In this letter we reconsider the FCNC processes in the littlest Higgs model with T-parity. We evaluate $d^j \rightarrow d^i \nu \bar{\nu}$ ($i \neq j$) amplitude and find disagreements with the previous results. In particular, we point out that there are no leftover singularities in the amplitude because the logarithmic divergent term in the Z -penguin and box diagrams is exactly canceled by contributions from additional interaction terms of heavy fermions and the Z boson. As a result, the FCNC amplitudes are determined by parameters in the original Lagrangian, and do not have to depend on unknown parameters of the ultraviolet completion of the theory.

The model we study in this letter is the littlest Higgs model with T-parity described

in detail in Ref. [10]. The gauge and the Higgs sector of the model is based on the $SU(5)/SO(5)$ nonlinear sigma model. The $SU(5)$ global symmetry is spontaneously broken down to $SO(5)$ by the scalar field Σ , which is transformed as **15** representation of the $SU(5)$. Σ is written as a 5×5 matrix and its $SU(5)$ transformation property is given as $\Sigma \rightarrow V\Sigma V^T$, where V is an arbitrary 5×5 unitary matrix with $\det V = 1$. The Nambu-Goldstone bosons are contained in a 5×5 matrix ξ , which is transformed under the $SU(5)$ as $\xi \rightarrow V\xi U$, where $U = U(V, \xi)$ is an $SO(5)$ matrix determined by V and ξ . $[SU(2) \times U(1)]^2$ subgroup of the $SU(5)$ is gauged and the $SU(2) \times U(1)$ electroweak gauge group of the Standard Model is assumed to be a diagonal subgroup of the $[SU(2) \times U(1)]^2$.

The vacuum expectation value of Σ is chosen as

$$\langle \Sigma \rangle = \xi_v \Sigma_0 \xi_v^T, \quad \xi_v = \langle \xi \rangle, \quad (1)$$

$$\Sigma_0 = \begin{pmatrix} 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 \end{pmatrix}, \quad (2)$$

$$\xi_v = \begin{pmatrix} 1 & 0 & 0 & 0 & 0 \\ 0 & \frac{c_v+1}{2} & \frac{is_v}{\sqrt{2}} & 0 & \frac{c_v-1}{2} \\ 0 & \frac{is_v}{\sqrt{2}} & c_v & 0 & \frac{is_v}{\sqrt{2}} \\ 0 & 0 & 0 & 1 & 0 \\ 0 & \frac{c_v-1}{2} & \frac{is_v}{\sqrt{2}} & 0 & \frac{c_v+1}{2} \end{pmatrix}, \quad (3)$$

where $c_v = \cos(v/(\sqrt{2}f))$ and $s_v = \sin(v/(\sqrt{2}f))$. $f = O(1 \text{ TeV})$ breaks $SU(5)$ down to $SO(5)$ and $[SU(2) \times U(1)]^2$ to $[SU(2) \times U(1)]_{\text{SM}}$. $v = (\sqrt{2}G_F)^{-1/2}$ is the electroweak symmetry breaking vacuum expectation value.

The gauge bosons of $SU(2)_i \times U(1)_i$ ($i = 1, 2$) are denoted as $W_i^{\pm,3}$ and B_i , respectively. T-odd combinations $W_H^{\pm,3} = (W_1^{\pm,3} - W_2^{\pm,3})/\sqrt{2}$ and $B_H = (B_1 - B_2)/\sqrt{2}$

receive masses of $O(f)$. T-even combinations $W_L^{\pm,3} = (W_1^{\pm,3} + W_2^{\pm,3})/\sqrt{2}$ and $B_L = (B_1 + B_2)/\sqrt{2}$ are identified as the Standard Model electroweak gauge bosons. After the electroweak symmetry breaking, the gauge fields of the mass eigenstates are given as

$$Z_L = W_L^3 \cos \theta_W - B_L \sin \theta_W, \quad A_L = W_L^3 \sin \theta_W + B_L \cos \theta_W, \quad (4)$$

$$Z_H = W_H^3 \cos \theta_H - B_H \sin \theta_H, \quad A_H = W_H^3 \sin \theta_H + B_H \cos \theta_H, \quad (5)$$

where A_L is the massless photon. θ_W is the Weinberg angle which is determined by the $[SU(2) \times U(1)]_{\text{SM}}$ gauge coupling constants g and g' as $\sin \theta_W = g'/\sqrt{g^2 + g'^2}$. The mixing angle of the T-odd gauge bosons is given by

$$\tan 2\theta_H = -\frac{gg'c_v^2s_v^2}{g^2 - g'^2/5 - (g^2 - g'^2)c_v^2s_v^2/2}. \quad (6)$$

Gauge boson masses are given as

$$m_{W_L}^2 = \frac{g^2 f^2}{2} s_v^2, \quad m_{Z_L}^2 = \frac{m_{W_L}^2}{\cos^2 \theta_W}, \quad (7)$$

$$m_{W_H}^2 = g^2 f^2 \left(1 - \frac{s_v^2}{2}\right), \quad (8)$$

$$\begin{aligned} m_{Z_H}^2 &= \frac{g^2 f^2}{c_H^2 - s_H^2} \left[\left(1 - \frac{c_v^2 s_v^2}{2}\right) c_H^2 - \frac{g'^2}{5g^2} \left(1 - \frac{5}{2} c_v^2 s_v^2\right) s_H^2 \right] \\ &= m_{W_H}^2 + O\left(\frac{v^4}{f^2}\right), \end{aligned} \quad (9)$$

$$\begin{aligned} m_{A_H}^2 &= \frac{g'^2 f^2}{5(c_H^2 - s_H^2)} \left[\left(1 - \frac{5}{2} c_v^2 s_v^2\right) c_H^2 - \frac{5g^2}{g'^2} \left(1 - \frac{c_v^2 s_v^2}{2}\right) s_H^2 \right] \\ &= \frac{g'^2 f^2}{5} \left(1 - \frac{5v^2}{4f^2} + O\left(\frac{v^4}{f^4}\right)\right), \end{aligned} \quad (10)$$

where $s_H = \sin \theta_H$ and $c_H = \cos \theta_H$.

In order to introduce the Standard Model fermions, the $SU(5) \supset [SU(2) \times U(1)]^2$ symmetry structure has to be extended because the $U(1)$ charges in the $SU(5)$ are not suitable to accommodate the hypercharge of the Standard Model fermions. This fact was implicitly assumed in Ref. [2]. A straightforward way is adding two $U(1)$ factors. Therefore, we consider the symmetry structure $SU(5) \times U(1)''_1 \times U(1)''_2 \supset [SU(2)_1 \times U(1)'_1 \times U(1)''_1] \times [SU(2)_2 \times U(1)'_2 \times U(1)''_2]$ hereafter. The $U(1)$ subgroups of the $SU(5)$ are renamed as $U(1)'_{1,2}$.

	$SU(2)_1$	$SU(2)_2$	Y'_1	Y'_2	Y''_1	Y''_2
$q_1 = \begin{pmatrix} u_1 \\ d_1 \end{pmatrix}$	2	1	$-\frac{3}{10}$	$-\frac{2}{10}$	$\frac{1}{3}$	$\frac{1}{3}$
$q_2 = \begin{pmatrix} u_2 \\ d_2 \end{pmatrix}$	1	2	$-\frac{2}{10}$	$-\frac{3}{10}$	$\frac{1}{3}$	$\frac{1}{3}$
t'_1	1	1	$\frac{2}{10}$	$-\frac{2}{10}$	$\frac{1}{3}$	$\frac{1}{3}$
t'_2	1	1	$-\frac{2}{10}$	$\frac{2}{10}$	$\frac{1}{3}$	$\frac{1}{3}$
$\ell_1 = \begin{pmatrix} \nu_1 \\ e_1 \end{pmatrix}$	2	1	$-\frac{3}{10}$	$-\frac{2}{10}$	0	0
$\ell_2 = \begin{pmatrix} \nu_2 \\ e_2 \end{pmatrix}$	1	2	$-\frac{2}{10}$	$-\frac{3}{10}$	0	0

Table 1: Quantum numbers of the left-handed fermion fields. Generation indices are suppressed.

The fermion sector of the littlest Higgs model with T-parity consists of three families of quark and lepton fields $q_{1,2}^k$, $\ell_{1,2}^k$, u_R^k , d_R^k , ν_R^k , e_R^k , q_{HR}^k and ℓ_{HR}^k with the generation index $k = 1, 2, 3$, and one set of the “top partner” fermions, $t'_{1,2}$ and $t'_{1R,2R}$. Quantum numbers of these fermion fields are summarized in Tables 1 and 2. $Y'_{1,2}$ and $Y''_{1,2}$ are charges of $U(1)'_{1,2}$ and $U(1)''_{1,2}$, respectively. Subgroups $U(1)_i \subset U(1)'_i \times U(1)''_i$ ($i = 1, 2$) with the charges $Y_i = Y'_i + Y''_i$ are gauged. The Standard Model hypercharge Y is given as $Y = Y_1 + Y_2$. “Mirror” fermions q_{HR} and ℓ_{HR} transform nonlinearly under $[SU(2)_1 \times U(1)'_1] \times [SU(2)_2 \times U(1)'_2]$. Transformation properties under T-parity are assigned as $(q_1, t'_1, t'_{1R}, \ell_1) \leftrightarrow (-q_2, -t'_2, -t'_{2R}, -\ell_2)$, $(u_R, d_R, \nu_R, e_R) \leftrightarrow (u_R, d_R, \nu_R, e_R)$ and $(q_{HR}, \ell_{HR}) \leftrightarrow (-q_{HR}, -\ell_{HR})$. From q_i , t'_i , t'_{iR} and ℓ_i ($i = 1, 2$), T-even fermion fields are constructed as $q_L = (q_1 - q_2)/\sqrt{2}$, $t'_+ = (t'_1 - t'_2)/\sqrt{2}$, $t'_{+R} = (t'_{1R} - t'_{2R})/\sqrt{2}$ and $\ell_L = (\ell_1 - \ell_2)/\sqrt{2}$, while T-odd ones are given as $q_H = (q_1 + q_2)/\sqrt{2}$, $t'_- = (t'_1 + t'_2)/\sqrt{2}$, $t'_{-R} = (t'_{1R} + t'_{2R})/\sqrt{2}$ and $\ell_H = (\ell_1 + \ell_2)/\sqrt{2}$. Note that $U(1)_{1,2}$ charges of the lepton

	$SU(2)_1$	$SU(2)_2$	Y'_1	Y'_2	Y''_1	Y''_2
u_R	1	1	0	0	$\frac{1}{3}$	$\frac{1}{3}$
d_R	1	1	0	0	$-\frac{1}{6}$	$-\frac{1}{6}$
$q_{HR} = \begin{pmatrix} u_{HR} \\ d_{HR} \end{pmatrix}$	—	—	—	—	$\frac{1}{3}$	$\frac{1}{3}$
t'_{1R}	1	1	$\frac{2}{10}$	$-\frac{2}{10}$	$\frac{1}{3}$	$\frac{1}{3}$
t'_{2R}	1	1	$-\frac{2}{10}$	$\frac{2}{10}$	$\frac{1}{3}$	$\frac{1}{3}$
ν_R	1	1	0	0	0	0
e_R	1	1	0	0	$-\frac{1}{2}$	$-\frac{1}{2}$
$\ell_{HR} = \begin{pmatrix} \nu_{HR} \\ e_{HR} \end{pmatrix}$	—	—	—	—	0	0

Table 2: Quantum numbers of the right-handed fermions. Generation indices are suppressed. q_{HR} and ℓ_{HR} transform nonlinearly under $[SU(2)_1 \times U(1)'_1] \times [SU(2)_2 \times U(1)'_2]$.

doublets $\ell_{1,2}$ are different from those used in Refs. [12, 10]. This charge assignment enables us to treat the lepton sector in the same way as the down-type quark sector.

$[SU(2)_1 \times U(1)'_1 \times U(1)''_1] \times [SU(2)_2 \times U(1)'_2 \times U(1)''_2]$ symmetric Yukawa coupling terms of the down-type quarks are written as

$$\mathcal{L}_{\text{down}} = \frac{i\lambda_d^{ij}f}{2\sqrt{2}} \sum_{p,q=1}^2 \sum_{x,y,z=3}^5 \epsilon_{pq}\epsilon_{xyz} \left[(\bar{\Psi}_2^{[5]i})_x (\Sigma)_{py} (\Sigma)_{qz} - (\bar{\Psi}_1^{[5]i}\Sigma_0)_x (\tilde{\Sigma})_{py} (\tilde{\Sigma})_{qz} \right] d_R^j + \text{H.c.}, \quad (11)$$

where λ_d^{ij} is the Yukawa coupling matrix and $\tilde{\Sigma}$ is the T-parity image of Σ . $\Psi_1^{[5]i}$ and $\Psi_2^{[5]i}$ are $SU(5) \times U(1)''_1 \times U(1)''_2$ multiplets with quantum numbers $(\mathbf{5}, -\frac{1}{6}, -\frac{1}{6})$ and $(\bar{\mathbf{5}}, -\frac{1}{6}, -\frac{1}{6})$, respectively. Quark doublets $q_{1,2}$ are embedded as

$$\Psi_1^{[5]} = \begin{pmatrix} q_1 \tilde{X}^\dagger \\ 0 \\ 0 \end{pmatrix}, \quad \Psi_2^{[5]} = \begin{pmatrix} 0 \\ 0 \\ q_2 X^\dagger \end{pmatrix}. \quad (12)$$

X and $\tilde{X}$ are $SU(2)_i$ singlet scalar fields with the $U(1)$ charges $(Y'_1, Y'_2, Y''_1, Y''_2) = (-\frac{4}{10}, -\frac{6}{10}, \frac{1}{2}, \frac{1}{2})$ and $(-\frac{6}{10}, -\frac{4}{10}, \frac{1}{2}, \frac{1}{2})$, respectively. Following Ref. [13], we replace X and $\tilde{X}$ with $(\Sigma_{33})^{-1/4}$ and its T-parity conjugate, respectively. Since the $U(1)_1 \times U(1)_2$ gauge charges of X are the same as those of $(\Sigma_{33})^{-1/4}$, this replacement maintains the gauge invariance of (11), whereas the $[U(1)'_1 \times U(1)''_1] \times [U(1)'_2 \times U(1)''_2]$ global symmetry is explicitly broken.

After the diagonalization of mass matrices, we have the following T-even (Dirac) fermions: three families of quarks and leptons u^k, d^k, ν^k and e^k ($k = 1, 2, 3$), and one top partner quark T_+ . Similarly, T-odd fermions are $u_H^k, d_H^k, \nu_H^k, e_H^k$, and T_- . T_+ and all the T-odd fermions have masses of $O(f)$. Flavor mixing among the T-even quarks is described by the CKM matrix V_{CKM} and an extra angle in the mixing of the top quark $t = u^3$ and T_+ . A parameter x_L ($0 < x_L < 1$) is introduced for this extra mixing angle [10]. The mixing matrix for the T-even lepton sector is the PMNS matrix U_{PMNS} . There are two independent mixing matrices in the T-odd gauge boson coupling with the fermions. We denote the mixing matrices at $\bar{d}_H^i \not{Z}_H (1 - \gamma_5) d^j$ and $\bar{\nu}_H^i \not{Z}_H (1 - \gamma_5) \nu^j$

couplings as $(V_{Hd})_{ij}$ and $(V_{H\nu})_{ij}$, respectively. Mixing matrices for other interaction terms are written in terms of V_{Hd} , $V_{H\nu}$, V_{CKM} and U_{PMNS} .

Main difference between our results and those of Ref. [10] originates from the gauge coupling of the Z_L boson and the “mirror” (right-handed and T-odd) fermions. The kinetic and gauge interaction terms of the mirror quarks are given as [12]

$$\mathcal{L}_{\text{kin}} = \frac{1}{2} \bar{\Psi}_R^{[5]i} \gamma^\mu \left(i\partial_\mu + g\hat{W}_\mu + g'\hat{B}_\mu^{\Psi'_R} \right) \Psi_R^{[5]i} + (\text{T-parity conjugate}). \quad (13)$$

$\Psi_R^{[5]i}$ is transformed as **5** representation under the $SU(5)$, and accommodate the mirror quarks as

$$\Psi_R^{[5]i} = \xi \begin{pmatrix} \tilde{\psi}_R^i \\ \chi_R^i \\ -i\sigma^2 q_{HR}^i \end{pmatrix}. \quad (14)$$

$\tilde{\psi}_R^i$ and χ_R^i are assumed to decouple from the effective theory. The gauge fields are written as

$$\begin{aligned} \hat{W} &= \frac{1}{2} \begin{pmatrix} W_L^3 & \sqrt{2}W_L^+ & 0 & 0 & 0 \\ \sqrt{2}W_L^- & -W_L^3 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & -W_L^3 & -\sqrt{2}W_L^- \\ 0 & 0 & 0 & -\sqrt{2}W_L^+ & W_L^3 \end{pmatrix} \\ &+ \frac{1}{2} \begin{pmatrix} W_H^3 & \sqrt{2}W_H^+ & 0 & 0 & 0 \\ \sqrt{2}W_H^- & -W_H^3 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & W_H^3 & \sqrt{2}W_H^- \\ 0 & 0 & 0 & \sqrt{2}W_H^+ & -W_H^3 \end{pmatrix}, \quad (15) \end{aligned}$$

$$\hat{B}^{\Psi'_R} = \frac{1}{6} \text{diag}(7, 7, 4, 1, 1)B_L + \frac{1}{10} \text{diag}(1, 1, -4, 1, 1)B_H. \quad (16)$$

We obtain the gauge interaction term of Z_L and W_L bosons with the mirror quarks as follows.

$$\mathcal{L}_{\text{gauge}} = g_Z \left[\left(\frac{1}{2} - \frac{2s_W^2}{3} + \delta_v \right) \bar{u}_{HR}^i \not{Z}_L u_{HR}^i + \left(-\frac{1}{2} + \frac{s_W^2}{3} \right) \bar{d}_{HR}^i \not{Z}_L d_{HR}^i \right]$$




Figure 1: Feynman diagrams for additional contribution to the Z_L penguin amplitudes in $d^j \rightarrow d^i$ transition. The box at the Z_L vertex denotes the correction term which is proportional to δ_v . ω^+ in (b) is the Nambu-Goldstone boson absorbed by W_H^+ .

$$+\frac{g}{\sqrt{2}}(1+\delta_v)\left(\bar{u}_{HR}^i W_L^+ d_{HR}^i + \bar{d}_{HR}^i W_L^- u_{HR}^i\right), \quad (17)$$

where $g_Z = \sqrt{g^2 + g'^2}$, $s_W = \sin \theta_W$ and $\delta_v = (c_v - 1)/2 = -v^2/(8f^2) + O(v^4/f^4)$. The terms proportional to δ_v are missing in the Feynman rules of Ref. [10]. These terms are generated because of the fact that the gauge field matrices $\hat{W}$ and $\hat{B}^{\Psi'_R}$ do not commute with ξ_v . The gauge interaction terms of the mirror leptons are derived in the same way. We find that there are the same correction terms that are proportional to δ_v in the interactions of the mirror neutrinos ν_{HR}^i .

The additional interaction terms that are proportional to δ_v in (17) affect the Z_L penguin contributions to the amplitudes for the $d^j \rightarrow d^i \nu \bar{\nu}$ FCNC processes. Relevant Feynman diagrams are shown in Fig. 1. We calculate all the relevant Z_L penguin and box diagrams with use of the 't Hooft–Feynman gauge and obtain the Wilson coefficients for $d^j \rightarrow d^i \nu \bar{\nu}$ as follows.

$$\mathcal{L}_{[d\nu]}^{\text{eff}} = C_{[d\nu]LL}^{ijlm}(\bar{d}^i \gamma^\mu \mathbf{L} d^j)(\bar{\nu}^l \gamma_\mu \mathbf{L} \nu^m), \quad (18)$$

$$C_{[d\nu]LL}^{ijlm} = -\frac{g^4}{(4\pi)^2 m_{W_L}^2} \left[\delta_{lm} \left(\sum_k \lambda_k X_{\text{SM}}(x_k) + \lambda_t \bar{X}_{\text{even}} \right) + \sum_{k,n} \lambda_n^{H\nu} \xi_k J^{\nu\bar{\nu}}(z_k, y_n) \right], \quad (19)$$

$$J^{\nu\bar{\nu}}(z_k, y_n) = \frac{v^2}{64f^2} \left[2z_k + z_k \log z_k + 3z_k \mathbf{f}_{0[1]}(z_k) - \mathbf{f}_3(z_k, y_n) + 7\mathbf{f}_2(z_k, y_n) \right]$$

$$-12\mathbf{g}_{2[1]}(z_k, y_n) - \frac{3r}{25}\mathbf{g}_{2[1]}(\frac{z_k}{r}, \frac{y_n}{r}) + \frac{6r}{5}\mathbf{g}_2(z_k, y_n, r) \Big], \quad (20)$$

where $\mathbf{L} = \frac{1}{2}(1 - \gamma_5)$, $x_k = m_{u^k}^2/m_{W_L}^2$, $z_k = m_{u^k}^2/m_{W_H}^2$, $y_n = m_{e_n}^2/m_{W_H}^2$, $r = m_{A_H}^2/m_{Z_H}^2$, $\lambda_k = (V_{\text{CKM}}^*)_{ki}(V_{\text{CKM}})_{kj}$, $\lambda_n^{H\nu} = (V_{H\nu}^*)_{nl}(V_{H\nu})_{nm}$, and $\xi_k = (V_{Hd}^*)_{ki}(V_{Hd})_{kj}$. Higher order terms in v/f expansions are neglected. We define the following classes of loop functions which are used in (20).

$$\mathbf{f}_n(y, z) = \frac{1}{y-z} \left(\frac{y^n \log y}{y-1} - \frac{z^n \log z}{z-1} \right), \quad (21a)$$

$$\mathbf{g}_n(x, y, z) = \frac{\mathbf{f}_n(x, z) - \mathbf{f}_n(y, z)}{x-y}, \quad (21b)$$

$$\mathbf{f}_{n[1]}(z) = \mathbf{f}_n(z, 1) = \frac{z^n \log z}{(z-1)^2} - \frac{1}{z-1}, \quad (21c)$$

$$\mathbf{g}_{n[1]}(x, y) = \mathbf{g}_n(x, y, 1) = \frac{\mathbf{f}_{n[1]}(x) - \mathbf{f}_{n[1]}(y)}{x-y}. \quad (21d)$$

For the T-even components of the Wilson coefficient (19), namely X_{SM} and $\bar{X}_{\text{even}}$, we obtain the same results as those in Ref. [10]. However, our result of the T-odd component, Eq. (20), differs from the corresponding formula of Ref. [10] in the following two points.

- There is no “leftover” $1/\epsilon$ singularity mentioned in [10]. The term proportional to δ_v in the $\bar{u}_{HR}\not{Z}_L u_{HR}$ coupling induces the following contribution to $J^{\nu\bar{\nu}}$ through the diagrams shown in Fig. 1.

$$\delta J^{\nu\bar{\nu}} = -\frac{v^2}{64f^2} z_k \left[\frac{1}{\epsilon} - \log \frac{m_{W_H}^2}{\mu^2} - \frac{1}{2} - \log z_k - 3\mathbf{f}_{0[1]}(z_k) \right]. \quad (22)$$

The $1/\epsilon$ term in (22) cancels the singularities from other contributions. For a crosscheck, we calculate $\delta J^{\nu\bar{\nu}}$ in the unitary gauge, with the diagram Fig. 1(a) only, and obtain the same result.

- The sign of the last term in the square brackets is opposite. This term is generated by the box diagrams which consists of d_H^k , ν_H^n , Z_H and A_H for the internal lines. The sign change is a consequence of the gauge charge assignments for the lepton doublets in Table 1.

In the same way, we obtain the corresponding quantity $J^{\mu\bar{\mu}}$, which appears in the $d^j \rightarrow d^i \ell^+ \ell^-$ Wilson coefficient, as follows.

$$J^{\mu\bar{\mu}}(z_k, y_n) = \frac{v^2}{64f^2} \left[2z_k + z_k \log z_k + 3z_k \mathbf{f}_{0[1]}(z_k) - \mathbf{f}_3(z_k, y_n) + 7\mathbf{f}_2(z_k, y_n) \right. \\ \left. + 6\mathbf{g}_{2[1]}(z_k, y_n) + \frac{3r}{25}\mathbf{g}_{2[1]}(\frac{z_k}{r}, \frac{y_n}{r}) + \frac{6r}{5}\mathbf{g}_2(z_k, y_n, r) \right]. \quad (23)$$

The difference from the formula in Ref. [10] is similar to the case of $J^{\nu\bar{\nu}}$.

In Ref. [11], the lepton flavor violating processes are studied. The formulae of the Wilson coefficients for the LFV effective operators are derived in a similar manner to those for the quark FCNC's in Ref. [10], so that the results depend on the ultraviolet singularities. However, these singularities are also canceled by the Z_L penguin contributions induced by the δ_v terms in the Z_L coupling with mirror neutrinos.

In Fig. 2, we show numerical values of $J^{\nu\bar{\nu}}$ and $J^{\mu\bar{\mu}}$ evaluated with use of Eq. (20) and (23), respectively. We take the mirror lepton mass $m_{eH} = 500$ GeV and $f = 1$ TeV for the calculation. We also show the values obtained by the formulae given in Ref. [10] (with $J_{\text{div}}^{\nu\bar{\nu}} = J_{\text{div}}^{\mu\bar{\mu}} = \frac{v^2}{64f^2} z \log \frac{(4\pi f)^2}{m_{WH}^2}$ and $J_{\text{div}}^{\nu\bar{\nu}} = J_{\text{div}}^{\mu\bar{\mu}} = 0$) in the same plot for comparison. The results with Eq. (20) and Eq. (23) look close to the results Ref. [10] with $J_{\text{div}}^{\nu\bar{\nu}} = J_{\text{div}}^{\mu\bar{\mu}} = 0$. Most of the numerical differences come from the contribution of $\delta J^{\nu\bar{\nu}}$. The effect of the sign change of the $d_H^k - \nu_H^n - Z_H - A_H$ box diagram is small.

Our result affects the predictions for $\Delta F = 1$ FCNC and LFV processes, in which Z_L penguin contributions of $O(v^2/f^2)$ are relevant. Here we present the results for the $K \rightarrow \pi \nu \bar{\nu}$ processes. In Fig. 3, the branching ratios of $K^+ \rightarrow \pi^+ \nu \bar{\nu}$ and $K_L \rightarrow \pi^0 \nu \bar{\nu}$ are shown as functions of the third generation T-odd quark mass $m_{H3} = m_{v_H^3} \simeq m_{d_H^3}$. As for other input parameters, we adopt the ‘‘Scenario 6’’ in Ref. [10]. In this scenario, it is assumed that the T-odd quark masses of the first and the second generations are equal and that the mixing parameter $\xi_3^{(K)} = (V_{Hd}^*)_{3s}(V_{Hd})_{3d}$ is pure imaginary. This parameter choice makes the T-odd contribution to the $K^0 - \bar{K}^0$ mixing negligibly small. We take the first and the second generation T-odd quark masses and all the T-odd lepton masses as 500 GeV. The parameters f and x_L are fixed as $f = 1$ TeV and $x_L = 0.5$. For




Figure 2: $J^{\nu\bar{\nu}}$ (solid lines) and $J^{\mu\bar{\mu}}$ (dashed lines) as functions of the mirror quark mass m_{uH} . Dark-grey/blue lines show the values calculated with Eqs. (20) and (23). Light-grey/magenta lines are the results of Ref. [10] with $J_{\text{div}}^{\nu\bar{\nu}} = J_{\text{div}}^{\mu\bar{\mu}} = \frac{v^2}{64f^2} z \log \frac{(4\pi f)^2}{m_{WH}^2}$ (thick lines) and $J_{\text{div}}^{\nu\bar{\nu}} = J_{\text{div}}^{\mu\bar{\mu}} = 0$ (thin lines).




Figure 3: Branching ratios of (a) $K^+ \rightarrow \pi^+ \nu \bar{\nu}$ and (b) $K_L \rightarrow \pi^0 \nu \bar{\nu}$ for $\text{Im} \xi_3 = 0.05$ (solid line) and $\text{Im} \xi_3 = 0.20$ (dashed line). Dark-grey/blue lines show the values calculated with Eq. (20). Light-grey/magenta lines are the results of Ref. [10] with $J_{\text{div}}^{\nu\bar{\nu}} = \frac{v^2}{64f^2} z \log \frac{(4\pi f)^2}{m_{W_H}^2}$. Horizontal solid lines show the Standard Model predictions. Dotted horizontal lines in (a) show 1σ range of the experimental value [14].

comparison, we also plot the branching ratios calculated by the formulae in Ref. [10] with $J_{\text{div}}^{\nu\bar{\nu}} = \frac{v^2}{64f^2} z \log \frac{(4\pi f)^2}{m_{W_H}^2}$. We see that both branching ratios can be significantly larger than the Standard Model prediction, although our values of the T-odd contributions are smaller than those in Ref. [10] in general.

In conclusion, we have revisited FCNC processes in the littlest Higgs model with T-parity. We have found that there is no divergence in the $d^i \rightarrow d^i \nu \bar{\nu}$ amplitude, that was reported earlier. This implies that FCNC processes can be insensitive to the physics at the cut-off scale. This is in contrast with the case of the littlest Higgs model without T-parity [15], where the leftover singularity exists. We have also re-evaluated the branching ratios of $K \rightarrow \pi \nu \bar{\nu}$ decays and found significant changes from previous results. We expect similar changes in other FCNC and LFV observables.

The work of T. G. and Y. O. is supported in part by the Grant-in-Aid for Science Research, Ministry of Education, Culture, Sports, Science and Technology, Japan, No. 16081211 and by the Grant-in-Aid for Science Research, Japan Society for the Promotion

References

- [1] N. Arkani-Hamed, A. G. Cohen and H. Georgi, Phys. Lett. B **513** (2001) 232 [arXiv:hep-ph/0105239]; N. Arkani-Hamed, A. G. Cohen, T. Gregoire and J. G. Wacker, JHEP **0208** (2002) 020 [arXiv:hep-ph/0202089]; N. Arkani-Hamed, A. G. Cohen, E. Katz, A. E. Nelson, T. Gregoire and J. G. Wacker, JHEP **0208** (2002) 021 [arXiv:hep-ph/0206020].
- [2] N. Arkani-Hamed, A. G. Cohen, E. Katz and A. E. Nelson, JHEP **0207** (2002) 034 [arXiv:hep-ph/0206021].
- [3] R. Barbieri and A. Strumia, Phys. Lett. B **462** (1999) 144 [arXiv:hep-ph/9905281].
- [4] C. Csaki, J. Hubisz, G. D. Kribs, P. Meade and J. Terning, Phys. Rev. D **67** (2003) 115002 [arXiv:hep-ph/0211124]. J. L. Hewett, F. J. Petriello and T. G. Rizzo, JHEP **0310** (2003) 062 [arXiv:hep-ph/0211218]. C. Csaki, J. Hubisz, G. D. Kribs, P. Meade and J. Terning, Phys. Rev. D **68** (2003) 035009 [arXiv:hep-ph/0303236].
- [5] H. C. Cheng and I. Low, JHEP **0309** (2003) 051 [arXiv:hep-ph/0308199]; H. C. Cheng and I. Low, JHEP **0408** (2004) 061 [arXiv:hep-ph/0405243].
- [6] N. Cabibbo, Phys. Rev. Lett. **10** (1963) 531; M. Kobayashi and T. Maskawa, Prog. Theor. Phys. **49** (1973) 652.
- [7] B. Pontecorvo, Sov. Phys. JETP **6**, 429 (1957) [Zh. Eksp. Teor. Fiz. **33**, 549 (1957)]; Sov. Phys. JETP **7**, 172 (1958) [Zh. Eksp. Teor. Fiz. **34**, 247 (1957)]; Sov. Phys. JETP **26**, 984 (1968) [Zh. Eksp. Teor. Fiz. **53**, 1717 (1967)]; Z. Maki, M. Nakagawa and S. Sakata, Prog. Theor. Phys. **28** (1962) 870.

- [8] J. Hubisz, S. J. Lee and G. Paz, JHEP **0606** (2006) 041 [arXiv:hep-ph/0512169];
M. Blanke, A. J. Buras, A. Poschenrieder, S. Recksiegel, C. Tarantino, S. Uhlig and
A. Weiler, Phys. Lett. B **646** (2007) 253 [arXiv:hep-ph/0609284].
- [9] M. Blanke, A. J. Buras, A. Poschenrieder, C. Tarantino, S. Uhlig and A. Weiler,
JHEP **0612** (2006) 003 [arXiv:hep-ph/0605214]; M. Blanke, A. J. Buras,
S. Recksiegel, C. Tarantino and S. Uhlig, Phys. Lett. B **657** (2007) 81
[arXiv:hep-ph/0703254]; M. Blanke, A. J. Buras, S. Recksiegel, C. Tarantino and
S. Uhlig, JHEP **0706** (2007) 082 [arXiv:0704.3329 [hep-ph]]; M. Blanke, A. J. Buras,
S. Recksiegel and C. Tarantino, arXiv:0805.4393 [hep-ph].
- [10] M. Blanke, A. J. Buras, A. Poschenrieder, S. Recksiegel, C. Tarantino, S. Uhlig and
A. Weiler, JHEP **0701** (2007) 066 [arXiv:hep-ph/0610298].
- [11] M. Blanke, A. J. Buras, B. Duling, A. Poschenrieder and C. Tarantino, JHEP **0705**
(2007) 013 [arXiv:hep-ph/0702136].
- [12] J. Hubisz and P. Meade, Phys. Rev. D **71** (2005) 035016 [arXiv:hep-ph/0411264].
- [13] C. R. Chen, K. Tobe and C. P. Yuan, Phys. Lett. B **640** (2006) 263
[arXiv:hep-ph/0602211]. A. Belyaev, C. R. Chen, K. Tobe and C. P. Yuan, Phys.
Rev. D **74** (2006) 115020 [arXiv:hep-ph/0609179].
- [14] V. V. Anisimovsky *et al.* [E949 Collaboration], Phys. Rev. Lett. **93** (2004) 031801
[arXiv:hep-ex/0403036].
- [15] A. J. Buras, A. Poschenrieder, S. Uhlig and W. A. Bardeen, JHEP **0611** (2006)
062 [arXiv:hep-ph/0607189].